

# EOS100850, a non-brain penetrant highly selective A<sub>2A</sub> receptor antagonist, uniquely maintains high potency within the adenosine rich tumor microenvironment


Houthuys E., Basilico P., Brouwer M., Deregnaucourt T., Detheux M., Driessens G., Gomes B., Leroy X., Marchante J., Marillier R., Swiercz J., Crosignani S.

AACR 2019 #3261

## SUMMARY


- High levels of extracellular adenosine drive tumor immunosuppression through A<sub>2A</sub> receptor signaling
- Adenosine concentrations in tumors are at least 10-fold higher compared to normal tissue (see posters #3278 and #4147)
- Clinical-stage competitor A<sub>2A</sub> receptor antagonists dramatically lose potency in a high adenosine environment
- EOS100850 is specifically designed for immuno-oncology:
  - Highly potent in high intratumoral adenosine concentrations
  - Potency not affected by adenosine levels
  - Long residence time on A<sub>2A</sub> receptor, resulting in prolonged PD effects *in vivo* (in vivo PD, see poster #4147)
  - Highly selective inhibitor of A<sub>2A</sub> receptor
  - Non-brain penetrant, avoiding potential CNS side-effects at doses needed to inhibit tumoral A<sub>2A</sub> receptors
  - Major metabolite is also a potent A<sub>2A</sub> receptor antagonist, and the potency is not affected by adenosine concentrations

## ADENOSINE-DRIVEN IMMUNOSUPPRESSION



Adapted from Ohta, Front Immunol 2016  
Antonic, Nat Rev Cancer 2013

## EOS100850 IS HIGHLY POTENT IN THE ADENOSINE RICH TUMOR MICROENVIRONMENT



**Fig. 1: EOS100850 is potent and outperforms clinical-stage competitors in conditions mimicking the tumor microenvironment.** Dose-response antagonistic activity of A<sub>2A</sub> receptor antagonists on human A<sub>2A</sub> receptor in tumor-like conditions. (A) HEK293-hA<sub>2A</sub>R cells were stimulated with 5 μM of adenosine in the presence of 2% (w/v) human serum albumin (HSA). Generation of cAMP was used as a readout. Antagonist activity is expressed as a percentage of the inhibition of adenosine activity. (B) Primary CD3<sup>+</sup> T cells were stimulated with a combination of anti-CD3 and anti-CD28 antibodies and 5 μM CGS-21680, in the presence of 50% human serum (HS). IL-2 was quantified by alphaLISA after a 72-hour incubation. The normalized data, expressed as percentage of IL-2 secretion rescue over vehicle control, are shown. Arcus Cpd I based on example 1 from patent application no. WO/2018/136700.


EOS100850 Assay	IC <sub>50</sub> in physiological environment	IC <sub>50</sub> in tumor-like environment
Potency in HEK-hA <sub>2A</sub> (cAMP)	2.24 nM	21.8 nM
Potency in primary human T cells (cAMP)	1.87 nM	1.83 nM
Potency in CD8 T cells in whole blood (pCREB)	-	11.2 nM

**Table 1. IC<sub>50</sub> values of EOS100850 on human A<sub>2A</sub> receptor in different assays mimicking physiological conditions and the tumor microenvironment.** EOS100850 effectively inhibits A<sub>2A</sub> receptor-mediated cAMP accumulation and pCREB formation, with IC<sub>50</sub> values in the low nanomolar range. EOS100850 potency is hardly affected by challenging conditions mimicking the tumor microenvironment (i.e. high plasma protein and high A<sub>2A</sub> receptor agonist conditions).

EOS100850 Assay	IC <sub>50</sub> in physiological environment	IC <sub>50</sub> in tumor-like environment
Potency in CHO-mA <sub>2A</sub> (cAMP)	12.9 nM	66.8 nM
Potency in primary mouse T cells (cAMP)	6.1 nM	3.13 nM
Potency in mouse whole blood (pCREB)	-	5.25 nM

**Table 2. IC<sub>50</sub> values of EOS100850 on mouse A<sub>2A</sub> receptor in different assays mimicking physiological conditions and the tumor microenvironment.**

## EOS100850 POTENCY IS NOT AFFECTED BY ADENOSINE LEVELS AND HAS A LONG RESIDENCE TIME ON A<sub>2A</sub> RECEPTOR



**Fig. 2: EOS100850 outperforms competitors in normal and adenosine-rich environments, and its potency is not affected by adenosine levels.** Human primary CD3<sup>+</sup> T cells were stimulated with different concentrations of adenosine (2% (w/v) HSA). cAMP was used as a readout. The potency of EOS100850 remains similar regardless of the adenosine concentration. As a result, EOS100850 is at least 100-1000 times more potent than competitors in the presence of 100 μM adenosine. Arcus Cpd I based on example 1 from patent application no. WO/2018/136700.

A <sub>2A</sub> R antagonist	K <sub>on</sub> (nM <sup>-1</sup> ·min <sup>-1</sup> )	K <sub>off</sub> (min <sup>-1</sup> )	RT (min)	K <sub>D</sub> (nM)
Rac-EOS100850	0.022 ± 0.010	0.006 ± 0.002	154 ± 48	0.30 ± 0.17
CPI-444	0.054 ± 0.003	0.045 ± 0.012	22 ± 6	0.83 ± 0.23
AZD4635	0.069 ± 0.029	0.181 ± 0.009	6 ± 1	2.6 ± 1.1


**Table 3: Receptor binding kinetics for rac-EOS100850 and competitor antagonists on hA<sub>2A</sub> receptor.** Association (K<sub>on</sub>) and dissociation rate constants (K<sub>off</sub>), residence times (RT), and (kinetic) K<sub>D</sub> values are shown. The long residence time on A<sub>2A</sub> receptor likely explains the 'insurmountable/non-competitive' profile of EOS100850, and results in a prolonged pharmacodynamic effect *in vivo* (see poster #4147).

## EOS100850 IS A HIGHLY SELECTIVE A<sub>2A</sub> RECEPTOR ANTAGONIST

Assay	EOS100850 IC <sub>50</sub>	Selectivity vs. A <sub>2A</sub>
A <sub>1</sub>	0.192 μM	270-fold
A <sub>2B</sub>	0.575 μM	1200-fold
A <sub>3</sub>	> 30 μM	>40000-fold

**Table 4. Activity of EOS100850 on human A<sub>1</sub>, A<sub>2B</sub>, and A<sub>3</sub> receptors.**


## EOS100850 IS NON-BRAIN PENETRANT



**Fig. 3: EOS100850 concentrations in mice following oral administration.** At T<sub>max</sub> as low as 1% of EOS100850 was detected in brain tissue compared to plasma. EOS100850 detected in the brain represented compound present in the residual blood in the collected brain tissue.

## EOS100850's MAJOR METABOLITE IS A POTENT A<sub>2A</sub> RECEPTOR ANTAGONIST

The major metabolite of EOS100850 (= 90%; EOS100612), identified both *in vitro* and *in vivo*, was synthesized and was found to be active on A<sub>2A</sub> receptors. Characterization in several assays revealed similar characteristics as the parent compound, i.e. high potency in tumor-like conditions, not affected by adenosine concentrations.



**Fig. 4: EOS100612 is potent in normal and adenosine-rich environments, and its potency is not affected by adenosine concentrations.** (A) Dose-response antagonistic activity of EOS100850 and EOS100612 on human A<sub>2A</sub> receptor in normal and tumor-like conditions. Generation of cAMP was used as a readout. Antagonist activity is expressed as a percentage of the inhibition of agonist activity. (B) Human primary CD3<sup>+</sup> T cells were stimulated with different concentrations of adenosine. cAMP was used as a readout. The potency of EOS100850 and EOS100612 remain similar regardless of the adenosine concentration.

EOS100612 Assay	IC <sub>50</sub> in physiological environment	IC <sub>50</sub> in tumor-like environment
Potency in HEK-hA <sub>2A</sub> (cAMP)	2.11 nM	73.2 nM
Potency in primary human T cells (cAMP)	2.40 nM	3.32 nM

**Table 5. IC<sub>50</sub> values of EOS100612, the principle metabolite of EOS100850, on human A<sub>2A</sub> receptor**

## CONCLUSIONS

EOS100850 is a novel, best-in-class A<sub>2A</sub> receptor antagonist designed for immuno-oncology

- Potent in high adenosine concentrations, a hallmark of the tumor microenvironment
- Potency is not affected by adenosine levels
- Long residence time on A<sub>2A</sub> receptor
- No CNS penetration and highly selective
- Major metabolite is highly potent A<sub>2A</sub> receptor antagonist

